ORIGINAL PAPER

UROLITHIASIS

The role of preoperative ureteral stenting in retrograde intrarenal surgery outcomes for renal stones: a matched-pair analysis

Samet Senel¹, Huseyin Gultekin¹, Yalcin Kizilkan¹, Cuneyt Ozden¹, Kazim Ceviz¹, Antonios Koudonas², Ahmet Halil Sevinc³

¹Department of Urology, Ankara City Hospital, Turkey

²First Department of Urology, School of Medicine, Aristotle University of Thessaloniki, Greece ³Department of Urology, Istanbul Kartal Dr. Lutfi Kirdar City Hospital, Istanbul, Turkey

Citation: Senel S, Gultekin H, Kizilkan Y, et al. The role of preoperative ureteral stenting in retrograde intrarenal surgery outcomes for renal stones: a matched-pair analysis. Cent European J Urol. 2024; 77: 668-673.

Article history

Submitted: Apr. 22, 2024 Accepted: Nov. 11, 2024 Published online: Dec. 17, 2024 Introduction Ureteral stenting is not routinely recommended, but it may be performed before or after retrograde intrarenal surgery (RIRS). We aimed to investigate the effect of preoperative ureteral stenting on the success rate and intraoperative, postoperative, and infectious complications in RIRS. **Material and methods** We retrospectively analysed the data of 581 patients who underwent RIRS. Demographic data, stone characteristics, presence of hydronephrosis, presence of congenital kidney anomaly and solitary kidney, duration of operation, and duration of hospitalisation were analysed. Intraoperative, postoperative, and infectious complications and the success rate of all operations were recorded. The patients were divided into 2 groups as prestented and non-prestented and matched in terms of age, sex, stone size, and number of stones. Ninety-four patients in the prestented group were matched with 282 patients in the non-prestented group with respect to age, sex, stone size, and number of stones (matched 1:3).

Corresponding author

Samet Senel Üniversiteler Mahallesi, 1604. Cadde No: 9 Çankaya, Ankara, Turkey samet_senel_umt@ hotmail.com **Results** The 2 groups were similar in terms of matching parameters and all other characteristics. After matching, the success rate was 77.7% (73/94) in the prestented group and 78% (220/282) in the non--prestented group, and there was no statistically significant difference between the 2 groups (p = 0.943). The intraoperative complication rate was statistically significantly higher in the non-prestented group (19.2% vs 28.7%, p = 0.046). Postoperative complications occurred in 22.3% of patients in the prestented group and 20.7% of patients in the non-prestented group (p = 0.429).

Conclusions Preoperative ureteral stenting in RIRS was not associated with the success rate or postoperative and infectious complications. However, preoperative stenting was effective in decreasing only grade 1 intraoperative complications.

Key Words: complication () preoperative ureteral stenting () RIRS () success

INTRODUCTION

Kidney stones comprise one of the most common urological diagnoses, and surgical treatments play an important role in the management of urinary stones. Regarding the surgical treatment of kidney stones, technological advancements contribute to a continuously evolving treatment algorithm, which at present includes the application of extracorporeal shockwave lithotripsy (ESWL), retrograde intrarenal surgery (RIRS), percutaneous nephrolithotomy (PCNL), and stone extraction procedures, which are performed either with open or minimally invasive approaches [1]. Among available modalities, RIRS demonstrates the most rapid, technology-driven evolvement, which is reflected in the expansion of its indication in the kidney stone surgical algorithm. Currently, the European Association of Urology (EAU) strongly recommends the application of RIRS for all patients with stones ≤ 2 cm, and even for larger stones in patients with contraindications for PCNL [2]. Similarly, the American Urological Association (AUA) recommends RIRS for every renal stone ≤ 2 cm, and additionally for patients who are not candidates for PCNL and bear larger stones [3].

Recent data from a multicentric database suggest that, under the current RIRS standards (methodology and indications), retrograde surgery for kidney stones is an effective modality for stone clearance, with only 11% of the patients undergoing additional RIRS sessions to achieve stone-free (SF) status [4]. At the same time, RIRS represents a safe endourological procedure; only 1.8% and 1.3% of the patients had complications > Clavien grade 2 in the intraoperative and postoperative setting, respectively [4]. Under the continuously increasing experience and familiarity with RIRS, several modifications are being tested with the intention of simplifying the whole process without compromising the results and the safety of the procedure. In the above patient cohort, 6.8% of the patients underwent a sheathless RIRS, while other reports describe the feasibility of performing effective and safe RIRS without the support of fluoroscopic guidance [5].

Another modification that is already applied by several urological centres is ureteral prestenting. Ureteral stents induce passive ureteral dilation in a predictable manner, but their advantage remains a topic of controversy among urologists. Both European and American guidelines recommend against routine prestenting in patients planned for RIRS on the basis of avoiding the cost of an additional procedure and the ureteral stent-related morbidity for achieving uncertain benefits [2, 3]. However, both the EAU and International Alliance of Urolithiasis recognise the possibility of improving SF rates and the safety of RIRS by placing a ureteral stent preoperatively [2, 6]. At present, there are no conclusive data relating to the decision for or against ureteral stenting before RIRS.

In the current study, we aimed to evaluate the effect of ureteral prestenting on the results and complications in patients who underwent RIRS in our clinic. To isolate the net effect of ureteral prestenting, special attention was paid to measuring surgical outcomes and complications in patient subcohorts, which were comparable in terms of every factor that may affect the results of RIRS in a parallel manner.

MATERIAL AND METHODS

Patients and clinical data

The data of 581 patients who underwent RIRS between January 2013 and May 2019 were obtained from the hospital information database retrospectively. The patients with prior urinary stone surgery, active urinary tract infection, concomitant ureteral stone, malignancy, history of radiation therapy, unsuccessful ureteral access sheath (UAS) placement because of ureteral stenosis or difficult ureter, and inadequate data were excluded from the study.

Demographic data of the patients (age, sex), stone characteristics (stone size, stone density, number of stones and lateralisation, stone location), presence of hydronephrosis, presence of congenital kidney anomaly and solitary kidney, duration of operation, and duration of hospitalisation were analysed. Intraoperative (modified Satava classification system), postoperative (modified Clavien-Dindo classification system), infectious (fever, urinary tract infection, sepsis and septic shock) complications, and success rates of all surgeries were recorded [7, 8].

All patients were diagnosed by non-contrast computed tomography (CT). Stone size was defined as the longest diameter of the stone and the sum of the longest diameters of all stones in the case of multiple stones.

We divided the patients into 2 groups: prestented and non-prestented. According to EAU guidelines, prestenting may improve the stone-free rate of ureteroscopic treatment of renal stones [2]. Prestenting was recommended to kidney stone patients before RIRS, and 1 or 2 weeks before RIRS a 4.8 F DJ stent was preoperatively placed in patients who accepted the presenting procedure. Among 581 patients, 94 patients were included in the presented group and 487 patients were included in the non-presented group. Ninety-four patients in the presented group were matched with patients in the non-prestented group with respect to age, sex, stone size, and number of stones. The 2 groups were compared in terms of intraoperative, postoperative, and infectious complications and characteristics of patients.

Surgical technique

All patients underwent RIRS under general anaesthesia in the lithotomy position. Ureterorenoscopy was performed with a 9.5 F semi-rigid ureterorenoscope (Karl Storz, Tuttingen, Germany) before RIRS. The DJ stent was extracted at the beginning of the procedure for the prestented group. After inserting the guidewire by semi-rigid ureterorenoscope, a 9.5-11 F UAS (Flexor® Uretheral access sheath, Cook Medical, USA) was inserted into the collecting system. A 7.5 F flexible ureterorenoscope (Karl Storz, Flex X2, GmbH, Tuttlingen, Germany) was placed into the collecting system through the access channel. The stones were fragmented using a holmium-yttrium-aluminium-garnet (Ho:YAG) laser (200–365 μ m) through the working channel of the flexible ureterorenoscope. At the end of the surgery, a 4.8 F DJ stent was placed into the collecting system. Postoperatively, the DJ stent was removed 2 weeks after RIRS, and all patients underwent non-contrast CT 4 weeks after RIRS to assess the success rate. Success was defined as the absence of any residual stones or fragments >2 mm in an asymptomatic patient. All procedures were performed by surgeons with at least 10 years of RIRS experience.

Statistical analysis

Data coding and statistical analyses were carried out on a computer using SPSS 22 software (IBM SPSS Statistics, IBM Corporation, Chicago, IL). Propensity score matching was performed using 4 preoperative (age, sex, stone size, and number of stones) variables by using predicted probability values in accordance with the nearest neighbour matching method in a 1:3 ratio. The conformity of the variables to the normal distribution was analysed using Shapiro-Wilk tests. Non-categorical parameters were presented as mean \pm standard deviation (SD) or median (minimum-maximum). The Mann-Whitney U test was used to compare non-categorical parameters, and χ^2 or Fisher's exact tests were used for categorical variables. A p-value <0.05 was accepted as statistically significant.

Bioethical standards

The institutional review board at Ankara City Hospital approved this study (approval number: E2.23.5862), and this study was prepared in accordance with the principles of the Declaration of Helsinki.

RESULTS

A total of 581 patients were included in the study. The mean age was 46.4 ± 14.5 years, and 366 (63%) patients were male. The median stone size was 16.2 ± 8.4 mm. Ninety-four (16.2%) patients were in the prestented group, and 487 (83.8%) patients were in the non-prestented group. There was no statistically significant difference between the 2 groups in terms of age (p = 0.309), sex (p = 0.113), stone size (p = 0.112), stone density (p = 0.39), number of stones (p = 0.844), stone location (p = 0.799), duration of operation (p = 0.702) and hospitalisation (p = 0.296), lateralisation (p = 0.549), presence of hydronephrosis (p = 0.381), congenital kidney anomaly (p = 0.484), and solitary kidney (p = 0.333). In the matching process, 94 patients in the prestented group were matched with 282 patients in the non-prestented group with respect to age, sex, stone size, and number of stones (matched 1:3). The two groups were similar in terms of matching parameters and all other characteristics (Table 1).

After matching, the success rate was 77.7% (73/94) in the prestented group and 78% (220/282) in the non-prestented group, and there was no statistically significant difference between the 2 groups (p = 0.943).

Eighteen (19.2%) patients had intraoperative complications (grade 1 in 9 patients, grade 2 in 9 patients) in the prestented group and 81 (28.7%) patients had intraoperative complications (grade 1 in 59 patients, grade 2 in 22 patients) in the nonprestented group. The intraoperative complication rate was statistically significantly higher in the non-prestented group (p = 0.046).

Postoperative complications occurred in 21 (22.3%) patients in the prestented group (grade 1 in 14 patients, grade 2 in 2 patients, grade 3 in 5 patients) and 57 (20.2%) patients in the non-prestented group (grade 1 in 47 patients, grade 2 in 2 patients, grade 3 in 8 patients), and the 2 groups were statistically similar in terms of postoperative complications (p = 0.429) (Table 2).

DISCUSSION

In the current study, we included 581 patients, who underwent a completed RIRS session. The surgical methodologies applied to the entire cohort were similar (ureteral sheath introduction, stone fragmentation with Ho:YAG laser, post-procedural ureteral stent placement), while a subgroup of the patient cohort was prestented to undergo RIRS with the ipsilateral ureter in a dilated state. Intraoperative and postoperative complications and success rate were evaluated in a standardised manner across the entire cohort. To minimise the effect of other factors that may interfere with the outcomes, we matched the prestented patients with non-prestented patients who had similar clinical characteristics. The comparison of the matched patient subgroups confirmed that they had no statistically significant

	Before matching			After matching		
Parameters	Prestented group (n = 94, 16.2%)	Non-prestented group (n = 487, 83.8%)	р	Prestented group (n = 94, 33.3%)	Non-prestented group (n = 282, 66.7%)	р
Age (years)	45.0 ±14.4	46.6 ±14.5	0.309*	45 ±14.4	45 ±14	0.931*
Sex Male, n (%) Female, n (%)	66 (70.2) 28 (29.8)	300 (61.6) 187 (38.4)	0.113**	66 (70.2) 28 (29.8)	199 (70.6) 83 (29.4)	0.948**
Stone size (mm), Mean ±SD	15.1 ±6.2	16.4 ±8.1	0.112*	15.1 ±6.2	15 ±6.2	0.193*
Stone density (HU), Mean ±SD	989.6 ±298.5	956.8 ±332.5	0.39*	989.6 ±298.5	941 ±339.6	0.179*
Number of stones Single, n (%) Multiple, n (%)	60 (63.8) 34 (36.2)	316 (64.9) 171 (35.1)	0.844**	60 (63.8) 34 (36.2)	179 (63.5) 103 (36.5)	0.951**
Stone location Pelvis, n (%) Upper, n (%) Middle, n (%) Lower, n (%) >1 calyx, n (%)	27 (28.7) 4 (4.3) 10 (10.6) 37 (39.4) 16 (17)	249 (42.9) 36 (6.2) 51 (8.8) 159 (27.4) 86 (14.8)	0.799**	27 (28.7) 4 (4.3) 10 (10.6) 37 (39.4) 16 (17)	123 (43.7) 19 (6.7) 31 (11) 79 (28) 30 (10.6)	0.519**
Duration of operation (min) (Mean ±SD)	50.2 ±15.7	55.1 ±18.7	0.702*	50.2 ±15.7	50.9 ±17.1	0.758*
Duration of hospitalisation (days) (Median) (min-max)	1 (1–17)	1 (1–75)	0.296*	1 (1–17)	1 (1–75)	0.921*
Lateralisation Right, n (%) Left, n (%)	42 (44.7) 52 (55.3)	234 (48) 253 (52)	0.549**	42 (44.7) 52 (55.3)	127 (45) 155 (55)	0.952**
Presence of hydronephrosis Yes, n (%) No, n (%)	59 (62.8) 35 (37.2)	282 (57.9) 205 (42.1)	0.381**	59 (62.8) 35 (37.2)	165 (58.5) 117 (41.5)	0.467**
Presence of congenital kidney anomaly Yes, n (%) No, n (%)	2 (2.1) 92 (97.9)	12 (2.5) 475 (97.5)	0.484**	2 (2.1) 92 (97.9)	12 (2.5) 475 (97.5)	0.602***
Presence of solitary kidney Yes, n (%) No, n (%)	5 (5.3) 89 (94.7)	16 (3.3) 471 (96.7)	0.333**	5 (5.3) 89 (94.7)	8 (2.8) 274 (97.2)	0.325***

 Table 1. Demographic, stone-related, and clinical characteristics of the prestented and non-prestented groups, and comparative analysis before and after matching

* Mann-Whitney U Test

** χ² test

*** Fisher's exact

HU – Hounsfield units; SD – standard deviation

difference in any clinical parameter. More importantly, ureteral prestenting did not affect stone clearance, because 78% of both groups were stone-free on week 4 postoperatively. Regarding intraoperative complications, prestented patients had significantly fewer events (19.2% vs 28.7% in the non-prestented group). This difference was more obvious in grade 1 complications, which manifested at a doubled rate in non-prestented patients. Postoperatively, both groups developed complication events at similar rates. In summary, the dilated ureter in the prestented patients was advantageous only in terms of minor intraoperative complications.

To compare our results with previous reports, we conducted a literature search and found that most studies demonstrate only slight or no superiority of ureteral prestenting. In 2018, Lee et al. [9] investigated the influence of the duration of ureteral prestenting on RIRS results and reported that prestenting did not affect RIRS outcomes, while it was beneficial only in reducing the necessity for intraoperative ureteral balloon dilation. However, Chen et al. [10] pooled the results of 11 studies and found that ureteral prestenting contributed significantly to stone clearance rates, while no difference was detected in terms of operative times and complication rates. Another report supporting ureteral prestenting was published by Giulioni et al. [11], who investigated the decisive factors determining stone clearance and complications in a cohort of 2,946 patients who underwent RIRS for lower pole stones. The report concluded that ureteral prestenting was among the factors independently affecting the stone clearance rate. In 2023, Assantachai et al. [12] compared

Table 2. Intraoperative, postoperative, and infectious complications and success rates of prestented and non-prestented groups in retrograde intrarenal surgery

	After matching				
Complications and success	Prestented group (n = 94, 16.2%)	Non-prestented group (n = 282, 66.7%)	р		
Intraoperative complication#					
None, n (%)	76 (80.9)	201 (71.3)	0.046*		
Grade 1, n (%) Minimal bleeding Minimal ureteral mucosal injury	9 (9.6) 5 4	59 (20.9) 33 26			
Grade 2, n (%) Bleeding requiring terminating surgery Ureteral mucosal injury requiring intervention	9 (9.6) 3 1	22 (7.8) 12 10			
Grade 3, n (%)	O (O)	0 (0)			
Postoperative non-infectious complication##					
None, n (%)	73 (77.7)	225 (79.8)	0.429*		
Grade 1, n (%) Haematuria Renal colic	14 (14.9) 10 4	47 (16.7) 29 18			
Grade 2, n (%) Blood transfusion	2 (2.1) 2	2 (0.7) 2			
Grade 3, n (%) Steinstrassen requiring ureterorenoscopy Perirenal abscess requiring drainage	5 (5.3) 4 1	8 (2.8) 7 1			
Grade 4, n (%)	O (O)	0 (0)			
Grade 5, n (%)	0 (0)	0 (0)			
Infectious complications None, n (%) Fever, n (%) Urinary tract infection, n (%) Sepsis, n (%) Septic shock, n (%)	87 (92.6) 3 (3.1) 2 (2.1) 1 (1.1) 1 (1.1)	257 (91.1) 16 (5.7) 8 (2.8) 0 (0) 1 (0.4)	0.327**		
Success Successful, n (%) Unsuccessful, n (%)	73 (77.7) 21 (22.3)	220 (78) 62 (22)	0.943*		

* χ² test

* Modified Satava Classification System

"Modified Clavien Classification System 2 patient groups that underwent RIRS and differed in the presence of a preoperative ureteral stent. The study reported no significant difference regarding stone clearance and complication rates. In line with the previous study, Jeong et al. [13] analysed the data of prestented vs non-prestented patients and concluded that ureteral prestenting was not superior in terms of stone clearance and complications. Regarding the paediatric population, Castellani et al. [14] compared prestented vs nonprestented patients and found that ureteral prestenting significantly increased the risk of postoperative infections, while no significant effect on stone clearance rates was detected.

In our opinion, the benefits of ureteral prestenting are limited and relate only to the placement of the UAS, while after the UAS introduction, there

was no difference in the following procedure, stone clearance, and complication rate. Indeed, UAS placement represents a demanding step of RIRS, and its uncomplicated success depends on anatomical and functional factors [15, 16]. In 2020, Sung et al. [17] demonstrated that ureteral prestenting significantly increased the success rate of UAS placement, while primary endpoints of RIRS were not affected. Similarly, Yuk et al. [18] reported that ureteral prestenting was advantageous only in terms of UAS placement success, and it should be considered in cases that are planned to be operated through UA. Recently, a meta-analysis by Law et al. [19] demonstrated the beneficial role of ureteral prestenting in UAS placement and additionally in stone clearance. The fact that ureteral prestenting seems to aid UAS placement renders

^{**} Fisher's Exact

questionable the benefit in patients who, on the basis of small stone size or surgeon's preference, are planned to undergo a sheathless RIRS.

This study has some limitations. The study was designed retrospectively, and therefore it is vulnerable to bias. Also, all surgeries were performed in a single centre. There is a lack of long-term follow-up results. In addition, data on which laser diameter was used in which case was not recorded in this retrospective study. However, we think that our study contributes to the literature on this subject, which includes conflicting results.

CONCLUSIONS

In this study, the data of prestented vs non-prestented patients who underwent RIRS after UAS placement were analysed, and no difference regarding stone clearance and postoperative complications were detected. Minor intraoperative complication rates were significantly lower in the prestented subgroup, which may be attributed to a more smooth UAS placement into the dilated ureter of these patients.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

FUNDING

This research received no external funding.

ETHICS APPROVAL STATEMENT

The study was approved by the institutional review board at Ankara City Hospital (approval number: E2.23.5862).

References

- Chung KJ, Kim JH, Min GE, et al. Changing Trends in the Treatment of Nephrolithiasis in the Real World. J Endourol. 2019; 33: 248-253.
- 2. EAU Guidelines. Edn. presented at the EAU Annual Congress Milan 2023.
- Assimos D, Krambeck A, Miller NL, et al. Surgical Management of Stones: American Urological Association/ Endourological Society Guideline, PART II. J Urol. 2016; 196: 1161-1169.
- Gauhar V, Chew BH, Traxer O, et al. Indications, preferences, global practice patterns and outcomes in retrograde intrarenal surgery (RIRS) for renal stones in adults: results from a multicenter database of 6669 patients of the global FLEXible ureteroscopy Outcomes Registry (FLEXOR). World J Urol. 2023; 41: 567-574.
- Laranjo Tinoco C, Coutinho A, Cardoso A, et al. Efficacy and safety of fluoroless ureteroscopy and retrograde intrarenal surgery for the treatment of urolithiasis: A comparative study. Actas Urol Esp (Engl Ed). 2023; 47: 535-542.
- Zeng G, Traxer O, Zhong W, et al. International Alliance of Urolithiasis guideline on retrograde intrarenal surgery. BJU Int. 2022; 131: 153-164.
- Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. Ann Surg. 2004; 240: 205-213.

- Satava RM. Identification and reduction of surgical error using simulation. Minim Invasive Ther Allied Technol. 2005; 14: 257-261.
- Lee MH, Lee IJ, Kim TJ, et al. The effect of short-term preoperative ureteral stenting on the outcomes of retrograde intrarenal surgery for renal stones. World J Urol. 2019; 37: 1435-1440.
- Chen H, Pan Y, Xiao M, et al. The Outcomes of Pre-Stenting on Renal and Ureteral Stones: A Meta-Analysis. Urol Int. 2021; 106: 495-503.
- Giulioni C, Castellani D, Somani BK, et al. The efficacy of retrograde intra-renal surgery (RIRS) for lower pole stones: results from 2946 patients. World J Urol. 2023; 41: 1407-1413.
- 12. Assantachai K, Srinualnad S, Leewansangtong S, Taweemonkongsap T, Liangkobkit K, Chotikawanich E. Surgical outcomes of patients who underwent retrograde intrarenal surgery using a ureteral access sheath to manage kidney stones sized 1-2 cm compared between patients who did and did not undergo preoperative ureteral stenting. Heliyon. 2023; 29; 9: e15801.
- Jeong JY, Cho KS, Jun DY, et al. Impact of Preoperative Ureteral Stenting in Retrograde Intrarenal Surgery for Urolithiasis. Medicina (Kaunas). 2023; 59: 744.
- 14. Castellani D, Somani BK, Ferretti S, et al. Role of Preoperative Ureteral Stent

on Outcomes of Retrograde Intra-Renal Surgery (RIRS) in Children. Results From a Comparative, Large, Multicenter Series. Urology. 2023; 173: 153-158.

- Mao L, Li Z, Wen J, Xu J, Zheng Z, Zhong W. Effect of bladder emptying status on the ureteral access sheath insertion resistance and following ureteral injury in RIRS: a prospective randomized controlled trial in academic hospital. World J Urol. 2023; 41: 2535-2540.
- Kim JK, Choi CI, Lee SH, et al. Silodosin for Prevention of Ureteral Injuries Resulting from Insertion of a Ureteral Access Sheath: A Randomized Controlled Trial. Eur Urol Focus. 2022; 8: 572-579.
- Sung LH, Cho DY. The role of preoperative ureteral stenting in retrograde intrarenal surgery in renal stone patients: a propensity score-matched study. Transl Androl Urol. 2020; 9: 276-283.
- Yuk HD, Park J, Cho SY, Sung LH, Jeong CW. The effect of preoperative ureteral stenting in retrograde Intrarenal surgery: a multicenter, propensity score-matched study. BMC Urol. 2020; 20: 147.
- Law YXT, Teoh JYC, Castellani D, et al. Role of pre-operative ureteral stent on outcomes of retrograde intra-renal surgery (RIRS): systematic review and meta-analysis of 3831 patients and comparison of Asian and non-Asian cohorts. World J Urol. 2022; 40: 1377-1389. ■